

Intro Physics Semester I**Name:**

Homework 15: Oscillation: One or two full answer questions will be marked. There will also be a mark for completeness. Homeworks are due usually the day after the chapter they are for is finished. Solutions will be posted soon thereafter. The solutions are intended to be (but not necessarily are) super-perfect and often go beyond a fully correct answer.

Answer Table**Name:**

	a	b	c	d	e		a	b	c	d	e
1.	<input type="radio"/>	31.	<input type="radio"/>								
2.	<input type="radio"/>	32.	<input type="radio"/>								
3.	<input type="radio"/>	33.	<input type="radio"/>								
4.	<input type="radio"/>	34.	<input type="radio"/>								
5.	<input type="radio"/>	35.	<input type="radio"/>								
6.	<input type="radio"/>	36.	<input type="radio"/>								
7.	<input type="radio"/>	37.	<input type="radio"/>								
8.	<input type="radio"/>	38.	<input type="radio"/>								
9.	<input type="radio"/>	39.	<input type="radio"/>								
10.	<input type="radio"/>	40.	<input type="radio"/>								
11.	<input type="radio"/>	41.	<input type="radio"/>								
12.	<input type="radio"/>	42.	<input type="radio"/>								
13.	<input type="radio"/>	43.	<input type="radio"/>								
14.	<input type="radio"/>	44.	<input type="radio"/>								
15.	<input type="radio"/>	45.	<input type="radio"/>								
16.	<input type="radio"/>	46.	<input type="radio"/>								
17.	<input type="radio"/>	47.	<input type="radio"/>								
18.	<input type="radio"/>	48.	<input type="radio"/>								
19.	<input type="radio"/>	49.	<input type="radio"/>								
20.	<input type="radio"/>	50.	<input type="radio"/>								
21.	<input type="radio"/>	51.	<input type="radio"/>								
22.	<input type="radio"/>	52.	<input type="radio"/>								
23.	<input type="radio"/>	53.	<input type="radio"/>								
24.	<input type="radio"/>	54.	<input type="radio"/>								
25.	<input type="radio"/>	55.	<input type="radio"/>								
26.	<input type="radio"/>	56.	<input type="radio"/>								
27.	<input type="radio"/>	57.	<input type="radio"/>								
28.	<input type="radio"/>	58.	<input type="radio"/>								
29.	<input type="radio"/>	59.	<input type="radio"/>								
30.	<input type="radio"/>	60.	<input type="radio"/>								

016 qmult 00100 1 4 5 easy deducto-memory: oscillation defined

1. “Let’s play *Jeopardy!* For \$100, the answer is: It is a motion that repeats itself in equal time periods: i.e., a periodic motion. Perfect repetition is an ideal case that is more or less closely approached in reality. The repeated motion is sometimes called a cycle.”

What is a/an _____, Alex?

- a) inhalation b) exhalation c) rotation d) acceleration e) oscillation

SUGGESTED ANSWER: (e)

Wrong answers:

- c) A rotation is not an oscillation, but oscillation can involve rotation in some cases.

Redaction: Jeffery, 2008jan01

016 qmult 00110 1 1 4 easy memory: period and frequency

2. Say P is the period of an oscillation. Say you observe N oscillations (or cycles) which take time NP , of course, The frequency of the oscillation (cycles per unit time) is given by:

- a) $f = NP$. b) $f = N/P$. c) $f = P/N$. d) $f = 1/P$. e) $f = N$.

SUGGESTED ANSWER: (d)

Behold:

$$f = \frac{N}{NP} = \frac{1}{P}.$$

Wrong answers:

- a) As Lurch would say AAAaaaargh.

Redaction: Jeffery, 2008jan01

016 qmult 00120 1 1 1 easy memory: unit of frequency

3. The MKS unit of frequency is the hertz (Hz). It is derived from the second:

- a) $1 \text{ Hz} = 1 \text{ s}^{-1}$. b) $1 \text{ Hz} = 1 \text{ s}$. c) $1 \text{ Hz} = 1 \text{ s}^{-2}$. d) $1 \text{ Hz} = 1 \text{ s}^2$. e) $1 \text{ Hz} = 1 \text{ s}^{-1/2}$.

SUGGESTED ANSWER: (a)

Wrong answers:

- a) As Lurch would say AAAaaaargh.

Redaction: Jeffery, 2008jan01

016 qmult 00200 1 1 2 easy memory: linear force law

4. The linear force law (AKA linear restoring force law, spring force law, Hooke’s law force, and simple harmonic oscillator force law) applies approximately to a wide variety of systems. In fact, almost any stable equilibrium system from molecules to bridges and beyond obeys the linear force law for sufficiently small perturbations of any component around the component’s stable equilibrium configuration. Components moving under the linear force law alone are in simple harmonic motion. Usually some damping force affects the system. The linear force law in one-dimension with the stable equilibrium point at the origin is given by:

- a) $F = -kx^{1/2}$. b) $F = -kx$. c) $F = -kx^{3/2}$. d) $F = -kx^2$. e) $F = kx$.

SUGGESTED ANSWER: (b)

Wrong answers:

- e) This gives the magnitude, but the sign is very wrong: this law gives an unstable equilibrium point at the origin.

Redaction: Jeffery, 2001jan01

016 qmult 00210 1 1 3 easy memory: linear force law ubiquity

5. To be in stable equilibrium, an object must be subject to a restoring force. The restoring force is zero at equilibrium and for any sufficiently small displacement from equilibrium tries to push the object back toward equilibrium. If the kinetic energy built up by the force pushing an object back to equilibrium is too large, then the object can **OVERSHOOT** the equilibrium and an oscillation results. Whether an oscillation results or not depends on the system. In real systems in the absence of any driver force, a damping force will usually dissipate mechanical energy above that of rest at the equilibrium point and cause the oscillation to damp out and the object to come to rest at the equilibrium point again. If the displacement is sufficiently small from a stable equilibrium, then the restoring force will be in most cases linear in the displacement: i.e., for a one-dimensional case

$$F = -kx ,$$

where k is a constant and x is the displacement from the equilibrium point. The restoring force in the small-displacement limit is often linear because:

- a) of no good reason. b) nature likes discontinuities. c) nature dislikes discontinuities.
d) nature is indifferent to discontinuities. e) it's lies, all lies.

SUGGESTED ANSWER: (c)

If the restoring force is continuous through the equilibrium point (which may be always true if one looks closely enough), then a Taylor's series expansion can be done at that point. Say $F(x)$ is the restoring force and $x = 0$ is the equilibrium. The Taylor's expansion is

$$F(x) = F(0) + x \left. \frac{dF}{dx} \right|_0 + \frac{x^2}{2} \left. \frac{d^2F}{dx^2} \right|_0 + \frac{x^3}{3} \left. \frac{d^3F}{dx^3} \right|_0 + \dots .$$

To be a restoring force $F(0) = 0$, $F(x < 0) > 0$, and $F(x > 0) < 0$. Thus, we see for small perturbations, the restoring force must be linear. Unless, $(dF/dx)|_0 = 0$. If $(dF/dx)|_0 = 0$, then we require $(d^2F/dx^2)|_0 = 0$ and $(d^3F/dx^3)|_0 < 0$. Such a 3rd-power restoring force seems to be very rare in nature—I know of no examples. One could design a system with a 3rd-power restoring force if one wanted to.

Wrong answers:

- e) A nonsense answer.

Redaction: Jeffery, 2008jan01

016 qmult 00220 1 1 3 easy memory: simple harmonic oscillator (SHO) 1

6. If the linear force alone acts on an object, the object is a simple harmonic oscillator (SHO) and undergoes simple harmonic motion (SHM). Apply Newton's 2nd law to 1-dimensional SHO with linear force $F = -kx$.

- a) $m \frac{d^2x}{dt^2} = kx$. b) $m \frac{dx}{dt} = -kx$. c) $m \frac{d^2x}{dt^2} = -kx$. d) $m \frac{d^2x}{dt^2} = -\frac{1}{kx}$.
e) $m \frac{d^2x}{dt^2} = -x$.

SUGGESTED ANSWER: (c)

Wrong answers:

- a) The minus sign is vital.

Redaction: Jeffery, 2008jan01

016 qmult 00230 1 1 1 easy memory: SHO solution 1

7. The general solution to the SHO differential equation

$$m \frac{d^2x}{dt^2} = -kx$$

is _____ where amplitude A and phase constant ϕ are set by initial conditions and angular frequency _____ is determined by the physics of the system itself. The motion of the solution is called simple harmonic motion (SHM).

- a) $x = A \cos(\omega t + \phi)$; $\omega = \sqrt{k/m}$ b) $x = A \cos(\omega t + \phi)$; $\omega = \sqrt{m/k}$
 c) $x = A(\omega t + \phi)$; $\omega = \sqrt{k/m}$ d) $x = A(\omega t + \phi)$; $\omega = \sqrt{m/k}$ e) $x = A(\omega t + \phi)$; $\omega = \sqrt{km}$

SUGGESTED ANSWER: (a)

For answer (a), we find

$$\text{LHS} = m(-\omega^2)x = -kx = \text{RHS}$$

which verifies that the answer is the solution for $\omega = \sqrt{k/m}$.

Wrong answers:

- b) Exactly wrong for omega.

Redaction: Jeffery, 2008jan01

016 qmult 00232 1 1 2 easy memory: SHO solution 2: SHO angular frequency formula

8. For a one-dimensional simple harmonic oscillator with mass m , force law $F = -kx$, and solution obeying $a = -\omega^2 x$, the formula for the angular frequency ω is given by:

- a) $\omega = m/k$. b) $\omega = \sqrt{k/m}$. c) $\omega = \sqrt{m/k}$. d) $\omega = k/m$. e) $\omega = \sqrt{km}$.

SUGGESTED ANSWER: (b)

From Newton's 2nd law we have

$$ma = -kx$$

and from the solution result we have

$$-m\omega^2 = -kx,$$

and so we find

$$\omega = \sqrt{\frac{k}{m}}.$$

Wrong answers:

- a) A nonsense answer.

Redaction: Jeffery, 2008jan01

016 qmult 00234 1 1 4 easy memory: SHO solution 3

9. A solution to Newton's 2nd law for the simple harmonic oscillator is simple harmonic motion—which is a melodious, uncomplicated motion. The solutions for displacement, velocity, and acceleration for a case when the motion of the oscillating objects starts at time $t = 0$ at its maximum displacement A are, respectively:

- a) $x = A \sin(\omega t)$; $v = \omega A \cos(\omega t)$; $x = -\omega^2 A \sin(\omega t)$.
 b) $x = A$; $v = \omega A$; $a = \omega^2 A$.
 c) $x = A \cos(\omega t)$; $v = -\omega A \sin(\omega t)$; $x = -\omega^2 A \cos(\omega t)$.
 d) $x = A \cos(\omega t)$; $v = -\omega A \cos(\omega t)$; $x = -\omega^2 A \cos(\omega t)$.
 e) $x = A \cos(\omega t)$; $v = \omega A \cos(\omega t)$; $x = \omega^2 A \cos(\omega t)$.

SUGGESTED ANSWER: (d)

Wrong answers:

- a) These are the solutions if it starts at $x = 0$ with positive velocity.
 b) These are just the maximum value solutions.

Redaction: Jeffery, 2008jan01

016 qmult 00240 1 1 1 easy memory: sinusoidal motion

10. In sinusoidal motion, an object's position as a function of time varies like a sine or:

- a) cosine curve. b) tangent curve. c) inverse tangent curve. d) inverse sine curve.
 e) straight line.

SUGGESTED ANSWER: (a)

Wrong answers:

b) Tangent curves have those infinities.

Redaction: Jeffery, 2008jan01

016 qmult 00250 1 1 3 easy memory: angular frequency

11. A sinusoid repeats its behavior every time its argument increases by 2π or 360° . An oscillation in time described by a sinusoid repeats in a time period called the period of the oscillation. The argument of the sinusoid in this case is $\omega t + \phi$, where ω is the angular frequency and ϕ is a phase constant set by initial conditions. For a period P , the angular frequency is given by:

a) $\omega = 2\pi P$. b) $\omega = P/(2\pi)$. c) $\omega = 2\pi/P$. d) $\omega = 1/(2\pi P)$. e) $\omega = 2\pi P^2$.

SUGGESTED ANSWER: (c)

Wrong answers:

a) Not dimensionally correct.

Redaction: Jeffery, 2008jan01

016 qmult 00260 1 1 3 easy memory: SHM time formulae related

12. For simple harmonic motion,

a) $f = P = \omega$. b) $P = 1/f = 2\pi/\omega$, $f = 1/P^2 = \omega/(2\pi)^2$, $\omega = 2\pi f = 2\pi/P$.
 c) $P = 1/f = 2\pi/\omega$, $f = 1/P = \omega/(2\pi)$, $\omega = 2\pi f = 2\pi/P$.
 d) $P = 1/f = 2\pi/\omega$, $f = 1/P = 2\pi\omega$, $\omega = 2\pi f = 2\pi/P$.
 e) $P = 1/f = 1/\omega$, $f = 1/P = \omega/(2\pi)$, $\omega = 2\pi f = 1/P$.

SUGGESTED ANSWER: (c)

Wrong answers:

a) A nonsense answer.

Redaction: Jeffery, 2008jan01

016 qmult 00270 3 4 1 hard deducto-memory: linearity of SHO equation

13. If $x_1(t)$ and $x_2(t)$ are solutions to a differential equation (DE), then the DE is linear if $c_1x_1(t) + c_2x_2(t)$ is also a solution where the c 's are arbitrary constants. The differential equation for the simple harmonic oscillator

$$\frac{d^2x}{dt^2} = -\omega^2x(t)$$

is:

- a) linear. b) non-linear. c) neither linear nor non-linear. d) both linear and non-linear.
 e) colinear.

SUGGESTED ANSWER: (a)

The SHO DE is linear by inspection.

Wrong answers:

- c) Not possible I think.
 d) Definitely not possible.
 e) Mischievous.

Redaction: Jeffery, 2001jan01

016 qmult 00280 1 1 2 easy memory: SHO energy

14. The mechanical energy of a 1-dimensional simple harmonic oscillator of mass m and force constant k is given by:

a) $E = kx_{\max}^2 = mv_{\max}^2$. b) $E = (1/2)kx_{\max}^2 = (1/2)mv_{\max}^2$. c) $E = (1/2)kx^2$.
 d) $E = (1/2)mv^2$. e) $E = mgy$.

SUGGESTED ANSWER: (b)

Wrong answers:

c) This is just the potential energy which is time varying.

d) This is just the kinetic energy which is time varying.

Redaction: Jeffery, 2008jan01

016 qmult 00290 1 1 3 easy memory: uniform circular motion and SHM

15. The projection of uniform circular motion on a line in the plane of rotation is:

- a) uniform circular motion. b) a Tusi-couple motion c) simple harmonic motion.
d) oblique motion. e) round motion.

SUGGESTED ANSWER: (c)

Wrong answers:

b) Well no, but Tusi-couple motion is related.

Fortran-95 Code

Redaction: Jeffery, 2008jan01

016 qmult 00300 1 1 4 easy memory: physical pendulum equation of motion

16. A physical pendulum is any rigid object held by a free pivot axis: the axis supports the pendulum against gravity, but exerts no torque about itself. The equation of motion of the physical pendulum (i.e., Newton's 2nd law applied to the physical pendulum) if only gravity torques it is:

$$\frac{d^2\theta}{dt^2} = -\frac{rmg}{I}\sin\theta ,$$

where θ is the angle a radius from the pivot axis to the center-of-mass axis makes with a downward vertical, r is the cylindrical radius from the pivot axis to the center-of-mass axis, m is the pendulum mass, and I is the pendulum rotational inertia about the pivot axis. This equation has no simple analytical solution because of:

- a) general considerations. b) the I factor. c) the r factor. d) the $\sin\theta$ factor.
e) darn good reasons.

SUGGESTED ANSWER: (d)

Wrong answers:

e) Not the best answer in this context.

Redaction: Jeffery, 2008jan01

016 qmult 00310 1 4 5 easy deducto-memory: small angle approximation

17. "Let's play *Jeopardy!* For \$100, the answer is:

$$\sin\theta \approx \theta ,$$

where θ is given in radians.

What is a/an _____, Alex?

- a) equality b) approximate equality c) inequality d) trigonometric function
e) small-angle approximation

SUGGESTED ANSWER: (e)

Wrong answers:

b) Not a best answer in this context.

Redaction: Jeffery, 2008jan01

016 qmult 00320 1 1 3 easy memory: physical pendulum in SHO approximation

18. The equation of motion of the physical pendulum (i.e., Newton's 2nd law applied to the physical pendulum) in the small-angle approximation is given by

$$\frac{d^2\theta}{dt^2} = -\frac{rmg}{I}\theta .$$

We recognize this equation as the differential equation of:

- a) Tusi-couple motion. b) uniform circular motion. c) simple harmonic motion.
 d) straight-line motion. e) uniform motion.

SUGGESTED ANSWER: (c)

Wrong answers:

- b) Nope.

Redaction: Jeffery, 2008jan01

016 qmult 00330 1 1 5 easy memory: simple pendulum period

19. The period of the physical pendulum (in the small-angle approximation) is

$$P = 2\pi\sqrt{\frac{I}{rmg}} .$$

If the pendulum is shrunk to a point-mass bob, we have the simple pendulum. In this case, _____ and _____.

- a) $I = (1/2)mr^2$; $P = 2\pi\sqrt{r/(2g)}$. b) $I = mr^2$; $P = 2\pi r/g$. c) $I = mr$; $P = 2\pi r/g$.
 d) $I = mr$; $P = 2\pi\sqrt{r/g}$. e) $I = mr^2$; $P = 2\pi\sqrt{r/g}$.

SUGGESTED ANSWER: (e)

Wrong answers:

- b) Wrong dimensionally.

Redaction: Jeffery, 2008jan01

016 qmult 00334 1 3 2 easy math: fiducial pendulum period

20. The period of a simple pendulum is given by

$$P = 2\pi\sqrt{\frac{r}{g}} ,$$

where r is the length of the pendulum and g is the gravitational field. A fiducial pendulum period (i.e., period that can be used as a standard for reference or quick estimation) is obtained for a pendulum of length exactly **ONE METER** and assuming that $g = 9.8 \text{ m/s}^2$ exactly. What is this fiducial period to 2 significant figures? To 4 significant figures?

- a) 1.0 s and 1.010 s. b) 2.0 s and 2.007 s. c) 2.0 s and 2.01 s. d) 1.0 s and 1.01 s.
 e) 1.0 s and 1.1 s.

SUGGESTED ANSWER: (b)

The rough calculation is

$$P = 2\pi\sqrt{\frac{r}{g}} \approx 6.3 \times \frac{1}{3.3} \approx 2 .$$

This result limits the possible right answers to answers (b) and (c). Since answer (c) has only 3 significant figures. Thus the right answer must be (b).

The more precise calculation to 4 significant figures is as follows:

$$P = 2\pi\sqrt{\frac{r}{g}} = 2.00708992 \dots \approx 2.007 .$$

Fortran Code

```
p=pi2*sqrt(1.d0/9.8d0)
print*,pi2,p
!      6.283185307179586 2.007089923154493
```

Wrong answers:

- c) This answer is only to 3 significant figures.

Redaction: Jeffery, 2001jan01

016 qmult 00336 2 3 5 moderate math: pendulum period sans calculator

21. Without using a calculator (you're on your honor here), what is approximately the period of a simple pendulum of length 9 m? **HINT:** Just look at what the simple pendulum period formula and reflect on the answer to the fiducial pendulum period question.

- a) 9.8 s. b) 162 s. c) 2.0 s. d) 1.0 s. e) 6.0 s.

SUGGESTED ANSWER: (e) For these modern, calculation-challenged students this is at least a moderate question.

Wrong answers:

- a) This is g and it looks suspiciously wrong.
 c) Whatever the answer is, it can't be the same as the fiducial pendulum period.
 d) Whatever the answer is, it can't be less than the fiducial pendulum period.

Redaction: Jeffery, 2001jan01

016 qmult 00340 3 5 4 tough thinking: pendulum period sans mass

22. Time for deep thought. If you are clever (and not like your physics professor) before doing any elaborate dynamical analysis, why should you know that the period of the simple pendulum is independent of mass?

- a) Mass is density times volume. Volume never comes into the simple pendulum problem. Ergo mass never comes into the simple pendulum problem.
 b) There no mass in the kinematic equations and thus mass can never affect the motion of anything. The simple pendulum is included in the set of anything.
 c) Physical intuition.
 d) For any particle acted on only by gravity and workless forces of constraint (e.g., normal forces such as those of a frictionless slope or of a frictionless wire for a sliding bead), the only force that can cause acceleration is the component of gravity in the direction allowed by the constraints. This component has mass as one of its factors and is the F in $F = ma$. Thus, the mass in 2nd Newton's law applied to the system cancels out from both sides of the equation. The acceleration is thus determined by g , the direction of gravity downward, and the directions allowed by the forces of constraint. In principle, the whole kinematics of the particle can be determined from knowing the acceleration for any location and the initial conditions. In practice, it might require elaborate calculations including computer calculations to find the whole kinematics. But mass does not come into the calculation. The simple pendulum fits the case just outlined. The ideal free massless pendulum arm exerts no force in the direction the pendulum bob can move. There is no air drag. The only force that can cause acceleration is a component of gravity. Consequently, the whole kinematics of the pendulum bob is independent of mass. The pendulum period is part of the kinematics. Therefore the pendulum period is independent of mass.
 e) Socrates has mass.
 All humans have mass.
 Ergo all humans are Socrates.

SUGGESTED ANSWER: (d)

Given that some answers are nonsense the students might be able to deduce the right answer. Also the question obeys the longest-answer-is-right rule.

Wrong answers:

- a) There is something wrong with this syllogism. Well first off in idealized physics we do have point masses and a point mass has no density or volume, and so the major premise is wrong for idealized physics. But even we grant that the major premise is true in reality and the minor premise is admittedly true, the conclusion still does not follow. We could still work the problem without specifying volume or density if only mass was needed to solve the dynamics.
 b) There is no mass in the kinematic equations, but there is acceleration. If acceleration depends on mass through $F = ma$, then the kinematics depends on mass. The fact that mass doesn't

- explicitly appear in the kinematic equations is not conclusive.
- c) Physical intuition is mostly just experience with a lot of problems. In this case, physical intuition will work *pour moi* because, I've done the simple pendulum problem many, many times. But it's not the best answer. Physical intuition is fallible.
 - e) Irrelevant even if valid. And it's probably not valid. I haven't noticed myself being Socrates ever—maybe Euripides now and then. If I'd ever taken a course in logic, I'd know the right terminology to point out the logical flaw. I suppose Socrates belongs to the set of mass-havers. So do all humans, but that doesn't imply all humans belong to the set Socrates. A Venn diagram would make it clear. My apologies to Woody Allen (1935–).

Redaction: Jeffery, 2001jan01

016 qmult 00350 1 5 5 easy thinking: pendulum clock and Galileo

23. You are Galileo Galilei (1564–1642) professor (untenured) of mathematics at the University of Pisa circa 1590. You are red-bearded and feisty (Italian word meaning . . .). Tired of people making fun of your redundant name (exactly so in Latin: Galileus Galileus) as if you were Humbert Humbert or some such and bored with dropping balls off the Leaning Tower in the piazza, you seek calm in the adjacent Cathedral of Pisa. There you notice that the Cathedral lanterns oscillate in the wind with a constant period no matter what the amplitude of the oscillation provided the amplitude isn't too large and that period only varies with lantern cord length and not lantern mass as far as you can tell. At once—you are an incomparable genius after all—you realize that pendulums would make great regulators for clocks because:

- a) it has **NEVER BEEN** been thought of before.
- b) it has **BEEN** thought of before.
- c) the hypnotic pendulum swinging motion would induce even deeper slumber in your less-amusing, clockwatching students.
- d) even a bad idea can make money. All that is needed is a great advertizing campaign.
- e) all clocks using pendulums as regulators for the motions of the hands and the energy input to keep the motions going would keep the periods of the hands very constant since small variations in amplitude that arise from somewhat irregular resistive and driver forces would have little effect on the pendulum period. Also the pendulum clocks could be kept synchronized to high accuracy despite varying amplitudes for the pendulum motion and masses for pendulum. Of course, the effective length of the pendulum does affect the period and has to be carefully adjusted for synchronization, but that is easy to do. It's one of those Eureka moments.

SUGGESTED ANSWER: (e)

Eliminating the nonsense answers leaves only one plausible answer. Of course this Galileo is fictional and the reasoning can only be very roughly along the lines of the real Galileo at some point in his life. He did experiment with pendulums and probably had the idea for a pendulum clock, but Christiaan Huygens (1629–1695) built the first real ones (Wikipedia 2007nov06).

Wrong answers:

- c) It probably would, but that's probably not what Galileo thought. I don't think that classrooms in the late 16th centuries had wall clocks. Possibly students relied on clock bells or chimes. We've mostly regressed from that level of sophistication. Where's our bell?
- d) The pendulum clock idea is not a bad idea.

Redaction: Jeffery, 2001jan01

016 qmult 00400 1 4 2 easy deducto-memory: damped oscillations

24. "Let's play *Jeopardy!* For \$100, the answer is: The motion of a system slowing down to rest at a stable equilibrium through a series of decaying oscillations about the equilibrium point."

What is _____, Alex?

- a) critically motion b) underdamped motion c) overdamped motion
- d) uniform circular motion e) all wet motion

SUGGESTED ANSWER: (b)

Wrong answers:

- a) No that's when the damping is just strong enough to stop the motion as fast as it can all else being equal.
- c) No that's when the damping is so strong that it slows the return to equilibrium relative to the critically damped case.
- e) If you chose this answer, then . . .

Redaction: Jeffery, 2001jan01

016 qmult 00530 1 4 4 easy deducto-memory: oscillatory resonance

25. "Let's play *Jeopardy!* For \$100, the answer is: The behavior of an oscillatory system driven at its natural or resonance frequency."

What is _____, Alex?

- a) underdamped oscillation
- b) simple harmonic motion
- c) uniform circular motion
- d) resonance or in resonance
- e) loco

SUGGESTED ANSWER: (d)

Wrong answers:

- e) Figuratively speaking maybe.

Redaction: Jeffery, 2001jan01

016 qmult 00540 1 4 1 easy deducto-memory: swing resonance of child

26. An oscillatory system driven at its resonance frequency will exhibit large oscillations. Every child—before they knew the name of torque—understood this when playing:

- a) on a swing.
- b) on a ladder.
- c) soccer.
- d) hopscotch.
- e) with matches.

SUGGESTED ANSWER: (a)

Wrong answers:

- e) You never did this, did you?

Redaction: Jeffery, 2001jan01

016 qfull 00210 1 3 0 easy math: SHO period, frequency, ω

27. There is a simple harmonic oscillator (SHO) that takes a time $\Delta t = 0.75$ s before it begins to repeat. What are its (a) period P , (b) frequency f (in hertz), and (c) angular frequency ω (in radians per second)?

SUGGESTED ANSWER:

Well (a) $P = 0.75$ s, (b) $f = 1/P = 4/3 = 1.33$ Hz, (c) $\omega = 2\pi f = 8.4$ rad/s.

Redaction: Jeffery, 2001jan01

016 qfull 00220 2 5 0 moderate thinking: maximum force of a SHO, etc.

28. A body of mass $m = 0.12$ kg is a simple harmonic oscillator (SHO) with equilibrium position $x = 0$, amplitude $x_{\max} = 8.5$ cm, period $P = 0.20$ s, and a linear (spring-like) restoring force.

- a) What is the force constant of the linear (or Hooke's law) force?
- b) What is the maximum absolute value of the force acting on the body?
- c) What is the maximum absolute value of the acceleration of the body?

SUGGESTED ANSWER:

- a) Well

$$\omega = \sqrt{\frac{k}{m}},$$

and so

$$P = 2\pi\sqrt{\frac{m}{k}}.$$

Thus

$$k = m\omega^2 = m \left(\frac{2\pi}{P} \right)^2 \approx 120 \text{ N/m}$$

to about 2-digit accuracy.

b) Behold:

$$|F|_{\max} = |kx_{\max}| = 10.5 \text{ N}$$

to about 2-digit accuracy. This is about 2 lb.

c) Behold:

$$|a|_{\max} = |\omega^2 x_{\max}| = \frac{|F|_{\max}}{m} \approx 85 \text{ m/s}^2$$

to about 2-digit accuracy.

Redaction: Jeffery, 2001jan01

016 qfull 00260 3 5 0 tough thinking: springs in parallel, oscillations

29. You have a block of mass m sandwiched between a bunch of springs in parallel. The whole system is a 1-dimensional system. The springs are attached to opposing walls. Some springs are from the left and some are from the right. The block sits on a level frictionless floor. The springs are ideal. Each spring has a force constant k_i and equilibrium position x_i for the center of the block: i.e., x_i is where the block center would be in equilibrium if only spring i were attached to the block.

- What is the expression for the net force on the mass?
- Derive the appropriate single-spring equivalent k (i.e., force constant) and x_{eq} (i.e., equilibrium position) expressions such that the net force expression changes to

$$F = -k(x - x_{\text{eq}}) .$$

Why is the x_{eq} the equilibrium position of the total system?

- Derive the expression for the total system ω in terms of the individual spring angular frequencies ω_i and the total system period P in terms of the individual spring periods P_i .

SUGGESTED ANSWER:

a) Behold:

$$F = - \sum_i k_i (x - x_i) .$$

It doesn't matter if the spring is attached from the left or the right: either way $(x - x_i) > 0$ gives a negative direction force and $(x - x_i) < 0$ gives a positive direction force.

b) Behold:

$$\begin{aligned} F &= - \sum_i k_i (x - x_i) \\ &= -x \sum_i k_i + \sum_i k_i x_i \\ &= -kx + k \frac{\sum_i k_i x_i}{k} \\ &= -kx + kx_{\text{eq}} \\ &= -k(x - x_{\text{eq}}) , \end{aligned}$$

where we define the equivalent single-spring equivalents k and x_{eq} by

$$k = \sum_i k_i \quad \text{and} \quad x_{\text{eq}} = \frac{\sum_i k_i x_i}{k} .$$

Note the equivalent k is just the sum of k_i and the equivalent equilibrium position is just the k_i weighted mean of the individual spring equilibrium positions. The equilibrium position total system is just x_{eq} since when $x = x_{\text{eq}}$ the net force is zero.

c) From 2nd law analysis, it follows immediately for

$$F = -k(x - x_{\text{eq}})$$

that the total system ω is $\omega = \sqrt{k/m}$, and thus

$$\omega = \sqrt{\frac{k}{m}} = \sqrt{\frac{\sum_i k_i}{m}} = \sqrt{\sum_i \omega_i^2}.$$

Now the total system period $P = 2\pi/\omega$ and $P_i = 2\pi/\omega_i$. Thus,

$$\frac{1}{P} = \sqrt{\sum_i \frac{1}{P_i^2}} \quad \text{or} \quad P = \frac{1}{\sqrt{\sum_i 1/P_i^2}}.$$

We have now pretty much solved the general N springs-in-parallel problem. Ah, the curse of generality as my old friend Francesco would say.

Redaction: Jeffery, 2001jan01

Equation Sheet for Introductory Physics Calculus-Based

This equation sheet is intended for students writing tests or reviewing material. Therefore it is neither intended to be complete nor completely explicit. There are fewer symbols than variables, and so some symbols must be used for different things: context must distinguish.

The equations are mnemonic. Students are expected to understand how to interpret and use them.

1 Constants

$$c = 2.99792458 \times 10^8 \text{ m/s} \approx 2.998 \times 10^8 \text{ m/s} \approx 3 \times 10^8 \text{ m/s} \approx 1 \text{ yr/yr} \approx 1 \text{ ft/ns} \quad \text{exact by definition}$$

$$e = 1.602176487(40) \times 10^{-19} \text{ C}$$

$$G = 6.67384(80) \times 10^{-11} \text{ N m}^2/\text{kg}^2 \quad (2012, \text{CODATA})$$

$$g = 9.8 \text{ m/s}^2 \quad \text{fiducial value}$$

$$k = \frac{1}{4\pi\epsilon_0} = 8.987551787 \dots \times 10^9 \approx 8.99 \times 10^9 \approx 10^{10} \text{ N m}^2/\text{C}^2 \text{ exact by definition}$$

$$k_{\text{Boltzmann}} = 1.3806504(24) \times 10^{-23} \text{ J/K} = 0.8617343(15) \times 10^{-4} \text{ eV/K} \approx 10^{-4} \text{ eV/K}$$

$$m_e = 9.10938215(45) \times 10^{-31} \text{ kg} = 0.510998910(13) \text{ MeV}$$

$$m_p = 1.672621637(83) \times 10^{-27} \text{ kg} = 938.272013(23), \text{ MeV}$$

$$\epsilon_0 = \frac{1}{\mu_0 c^2} = 8.8541878176 \dots \times 10^{-12} \text{ C}^2/(\text{N m}^2) \approx 10^{-11} \quad \text{vacuum permittivity (exact by definition)}$$

$$\mu_0 = 4\pi \times 10^{-7} \text{ N/A}^2 \quad \text{exact by definition}$$

2 Geometrical Formulae

$$C_{\text{cir}} = 2\pi r \quad A_{\text{cir}} = \pi r^2 \quad A_{\text{sph}} = 4\pi r^2 \quad V_{\text{sph}} = \frac{4}{3}\pi r^3$$

$$\Omega_{\text{sphere}} = 4\pi \quad d\Omega = \sin\theta \, d\theta \, d\phi$$

3 Trigonometry Formulae

$$\frac{x}{r} = \cos\theta \quad \frac{y}{r} = \sin\theta \quad \frac{y}{x} = \tan\theta = \frac{\sin\theta}{\cos\theta} \quad \cos^2\theta + \sin^2\theta = 1$$

$$\csc\theta = \frac{1}{\sin\theta} \quad \sec\theta = \frac{1}{\cos\theta} \quad \cot\theta = \frac{1}{\tan\theta}$$

$$c^2 = a^2 + b^2 \quad c = \sqrt{a^2 + b^2 - 2ab\cos\theta_c} \quad \frac{\sin\theta_a}{a} = \frac{\sin\theta_b}{b} = \frac{\sin\theta_c}{c}$$

$$f(\theta) = f(\theta + 360^\circ)$$

$$\sin(\theta + 180^\circ) = -\sin(\theta) \quad \cos(\theta + 180^\circ) = -\cos(\theta) \quad \tan(\theta + 180^\circ) = \tan(\theta)$$

$$\sin(-\theta) = -\sin(\theta) \quad \cos(-\theta) = \cos(\theta) \quad \tan(-\theta) = -\tan(\theta)$$

$$\sin(\theta + 90^\circ) = \cos(\theta) \quad \cos(\theta + 90^\circ) = -\sin(\theta) \quad \tan(\theta + 90^\circ) = -\tan(\theta)$$

$$\sin(180^\circ - \theta) = \sin(\theta) \quad \cos(180^\circ - \theta) = -\cos(\theta) \quad \tan(180^\circ - \theta) = -\tan(\theta)$$

$$\sin(90^\circ - \theta) = \cos(\theta) \quad \cos(90^\circ - \theta) = \sin(\theta) \quad \tan(90^\circ - \theta) = \frac{1}{\tan(\theta)} = \cot(\theta)$$

$$\sin(a + b) = \sin(a)\cos(b) + \cos(a)\sin(b) \quad \cos(a + b) = \cos(a)\cos(b) - \sin(a)\sin(b)$$

$$\sin(2a) = 2\sin(a)\cos(a) \quad \cos(2a) = \cos^2(a) - \sin^2(a)$$

$$\sin(a)\sin(b) = \frac{1}{2}[\cos(a - b) - \cos(a + b)] \quad \cos(a)\cos(b) = \frac{1}{2}[\cos(a - b) + \cos(a + b)]$$

$$\sin(a)\cos(b) = \frac{1}{2}[\sin(a - b) + \sin(a + b)]$$

$$\sin^2\theta = \frac{1}{2}[1 - \cos(2\theta)] \quad \cos^2\theta = \frac{1}{2}[1 + \cos(2\theta)] \quad \sin(a)\cos(a) = \frac{1}{2}\sin(2a)$$

$$\cos(x) - \cos(y) = -2\sin\left(\frac{x+y}{2}\right)\sin\left(\frac{x-y}{2}\right)$$

$$\cos(x) + \cos(y) = 2\cos\left(\frac{x+y}{2}\right)\cos\left(\frac{x-y}{2}\right)$$

$$\sin(x) + \sin(y) = 2\sin\left(\frac{x+y}{2}\right)\cos\left(\frac{x-y}{2}\right)$$

4 Approximation Formulae

$$\frac{\Delta f}{\Delta x} \approx \frac{df}{dx} \quad \frac{1}{1-x} \approx 1+x : (x \ll 1)$$

$$\sin\theta \approx \theta \quad \tan\theta \approx \theta \quad \cos\theta \approx 1 - \frac{1}{2}\theta^2 \quad \text{all for } \theta \ll 1$$

5 Quadratic Formula

$$\text{If } 0 = ax^2 + bx + c, \quad \text{then } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = -\frac{b}{2a} \pm \sqrt{\left(\frac{b}{2a}\right)^2 - \frac{c}{a}}$$

6 Vector Formulae

$$a = |\vec{a}| = \sqrt{a_x^2 + a_y^2} \quad \theta = \tan^{-1} \left(\frac{a_y}{a_x} \right) + \pi? \quad \vec{a} + \vec{b} = (a_x + b_x, a_y + b_y)$$

$$a = |\vec{a}| = \sqrt{a_x^2 + a_y^2 + a_z^2} \quad \phi = \tan^{-1} \left(\frac{a_y}{a_x} \right) + \pi? \quad \theta = \cos^{-1} \left(\frac{a_z}{a} \right)$$

$$\vec{a} + \vec{b} = (a_x + b_x, a_y + b_y, a_z + b_z)$$

$$\vec{a} \cdot \vec{b} = ab \cos \theta = a_x b_x + a_y b_y + a_z b_z$$

$$\vec{c} = \vec{a} \times \vec{b} = ab \sin(\theta) \hat{c} = (a_y b_z - b_y a_z, a_z b_x - b_z a_x, a_x b_y - b_x a_y)$$

7 Differentiation and Integration Formulae

$$\frac{d(x^p)}{dx} = px^{p-1} \quad \text{except for } p = 0; \quad \frac{d(x^0)}{dx} = 0 \quad \frac{d(\ln|x|)}{dx} = \frac{1}{x}$$

Taylor's series

$$f(x) = \sum_{n=0}^{\infty} \frac{(x-x_0)^n}{n!} f^{(n)}(x_0)$$

$$= f(x_0) + (x-x_0)f^{(1)}(x_0) + \frac{(x-x_0)^2}{2!}f^{(2)}(x_0) + \frac{(x-x_0)^3}{3!}f^{(3)}(x_0) + \dots$$

$$\int_a^b f(x) dx = F(x)|_a^b = F(b) - F(a) \quad \text{where} \quad \frac{dF(x)}{dx} = f(x)$$

$$\int x^n dx = \frac{x^{n+1}}{n+1} \quad \text{except for } n = -1; \quad \int \frac{1}{x} dx = \ln|x|$$

8 One-Dimensional Kinematics

$$v_{\text{avg}} = \frac{\Delta x}{\Delta t} \quad v = \frac{dx}{dt} \quad a_{\text{avg}} = \frac{\Delta v}{\Delta t} \quad a = \frac{dv}{dt} = \frac{d^2x}{dt^2}$$

$$v = at + v_0 \quad x = \frac{1}{2}at^2 + v_0t + x_0 \quad v^2 = v_0^2 + 2a(x - x_0)$$

$$x = \frac{1}{2}(v_0 + v)t + x_0 \quad x = -\frac{1}{2}at^2 + vt + x_0 \quad g = 9.8 \text{ m/s}^2$$

$$x_{\text{rel}} = x_2 - x_1 \quad v_{\text{rel}} = v_2 - v_1 \quad a_{\text{rel}} = a_2 - a_1$$

$$x' = x - v_{\text{frame}}t \quad v' = v - v_{\text{frame}} \quad a' = a$$

9 Two- and Three-Dimensional Kinematics: General

$$\vec{v}_{\text{avg}} = \frac{\Delta \vec{r}}{\Delta t} \quad \vec{v} = \frac{d\vec{r}}{dt} \quad \vec{a}_{\text{avg}} = \frac{\Delta \vec{v}}{\Delta t} \quad \vec{a} = \frac{d\vec{v}}{dt} = \frac{d^2\vec{r}}{dt^2}$$

10 Projectile Motion

$$x = v_{x,0}t \quad y = -\frac{1}{2}gt^2 + v_{y,0}t + y_0 \quad v_{x,0} = v_0 \cos \theta \quad v_{y,0} = v_0 \sin \theta$$

$$t = \frac{x}{v_{x,0}} = \frac{x}{v_0 \cos \theta} \quad y = y_0 + x \tan \theta - \frac{x^2 g}{2v_0^2 \cos^2 \theta}$$

$$x_{\text{for } y \text{ max}} = \frac{v_0^2 \sin \theta \cos \theta}{g} \quad y_{\text{max}} = y_0 + \frac{v_0^2 \sin^2 \theta}{2g}$$

$$x(y = y_0) = \frac{2v_0^2 \sin \theta \cos \theta}{g} = \frac{v_0^2 \sin(2\theta)}{g} \quad \theta_{\text{for max}} = \frac{\pi}{4} \quad x_{\text{max}}(y = y_0) = \frac{v_0^2}{g}$$

$$x(\theta = 0) = \pm v_0 \sqrt{\frac{2(y_0 - y)}{g}} \quad t(\theta = 0) = \sqrt{\frac{2(y_0 - y)}{g}}$$

11 Relative Motion

$$\vec{r} = \vec{r}_2 - \vec{r}_1 \quad \vec{v} = \vec{v}_2 - \vec{v}_1 \quad \vec{a} = \vec{a}_2 - \vec{a}_1$$

12 Polar Coordinate Motion and Uniform Circular Motion

$$\omega = \frac{d\theta}{dt} \quad \alpha = \frac{d\omega}{dt} = \frac{d^2\theta}{dt^2}$$

$$\vec{r} = r\hat{r} \quad \vec{v} = \frac{d\vec{r}}{dt} = \frac{dr}{dt}\hat{r} + r\omega\hat{\theta} \quad \vec{a} = \frac{d^2\vec{r}}{dt^2} = \left(\frac{d^2r}{dt^2} - r\omega^2\right)\hat{r} + \left(r\alpha + 2\frac{dr}{dt}\omega\right)\hat{\theta}$$

$$\vec{v} = r\omega\hat{\theta} \quad v = r\omega \quad a_{\text{tan}} = r\alpha$$

$$\vec{a}_{\text{centripetal}} = -\frac{v^2}{r}\hat{r} = -r\omega^2\hat{r} \quad a_{\text{centripetal}} = \frac{v^2}{r} = r\omega^2 = v\omega$$

13 Very Basic Newtonian Physics

$$\vec{r}_{\text{cm}} = \frac{\sum_i m_i \vec{r}_i}{m_{\text{total}}} = \frac{\sum_{\text{sub}} m_{\text{sub}} \vec{r}_{\text{cm sub}}}{m_{\text{total}}} \quad \vec{v}_{\text{cm}} = \frac{\sum_i m_i \vec{v}_i}{m_{\text{total}}} \quad \vec{a}_{\text{cm}} = \frac{\sum_i m_i \vec{a}_i}{m_{\text{total}}}$$

$$\vec{r}_{\text{cm}} = \frac{\int_V \rho(\vec{r}) \vec{r} dV}{m_{\text{total}}}$$

$$\vec{F}_{\text{net}} = m\vec{a} \quad \vec{F}_{21} = -\vec{F}_{12} \quad F_g = mg \quad g = 9.8 \text{ m/s}^2$$

$$\vec{F}_{\text{normal}} = -\vec{F}_{\text{applied}} \quad F_{\text{linear}} = -kx$$

$$f_{\text{normal}} = \frac{T}{r} \quad T = T_0 - F_{\text{parallel}}(s) \quad T = T_0$$

$$F_{\text{f static}} = \min(F_{\text{applied}}, F_{\text{f static max}}) \quad F_{\text{f static max}} = \mu_{\text{static}} F_{\text{N}} \quad F_{\text{f kinetic}} = \mu_{\text{kinetic}} F_{\text{N}}$$

$$v_{\text{tangential}} = r\omega = r \frac{d\theta}{dt} \quad a_{\text{tangential}} = r\alpha = r \frac{d\omega}{dt} = r \frac{d^2\theta}{dt^2}$$

$$\vec{a}_{\text{centripetal}} = -\frac{v^2}{r} \hat{r} \quad \vec{F}_{\text{centripetal}} = -m \frac{v^2}{r} \hat{r}$$

$$F_{\text{drag, lin}} = bv \quad v_{\text{T}} = \frac{mg}{b} \quad \tau = \frac{v_{\text{T}}}{g} = \frac{m}{b} \quad v = v_{\text{T}}(1 - e^{-t/\tau})$$

$$F_{\text{drag, quad}} = bv^2 = \frac{1}{2} C \rho A v^2 \quad v_{\text{T}} = \sqrt{\frac{mg}{b}}$$

14 Energy and Work

$$dW = \vec{F} \cdot d\vec{s} \quad W = \int \vec{F} \cdot d\vec{s} \quad KE = \frac{1}{2}mv^2 \quad E_{\text{mechanical}} = KE + PE$$

$$P_{\text{avg}} = \frac{\Delta W}{\Delta t} \quad P = \frac{dW}{dt} \quad P = \vec{F} \cdot \vec{v}$$

$$\Delta KE = W_{\text{net}} \quad \Delta PE_{\text{of a conservative force}} = -W_{\text{by a conservative force}} \quad \Delta E = W_{\text{nonconservative}}$$

$$F = -\frac{dPE}{dx} \quad \vec{F} = -\nabla PE \quad PE = \frac{1}{2}kx^2 \quad PE = mgy$$

15 Momentum

$$\vec{F}_{\text{net}} = m\vec{a}_{\text{cm}} \quad \Delta KE_{\text{cm}} = W_{\text{net,external}} \quad \Delta E_{\text{cm}} = W_{\text{not}}$$

$$\vec{p} = m\vec{v} \quad \vec{F}_{\text{net}} = \frac{d\vec{p}}{dt} \quad \vec{F}_{\text{net}} = \frac{d\vec{p}_{\text{total}}}{dt}$$

$$m\vec{a}_{\text{cm}} = \vec{F}_{\text{net non-flux}} + (\vec{v}_{\text{flux}} - \vec{v}_{\text{cm}}) \frac{dm}{dt} = \vec{F}_{\text{net non-flux}} + \vec{v}_{\text{rel}} \frac{dm}{dt}$$

$$v = v_0 + v_{\text{ex}} \ln\left(\frac{m_0}{m}\right) \quad \text{rocket in free space}$$

16 Collisions

$$\vec{I} = \int_{\Delta t} \vec{F}(t) dt \quad \vec{F}_{\text{avg}} = \frac{\vec{I}}{\Delta t} \quad \Delta p = \vec{I}_{\text{net}}$$

$$\vec{p}_{1i} + \vec{p}_{2i} = \vec{p}_{1f} + \vec{p}_{2f} \quad \vec{v}_{\text{cm}} = \frac{\vec{p}_1 + \vec{p}_2}{m_{\text{total}}}$$

$$KE_{\text{total } f} = KE_{\text{total } i} \quad \text{1-d Elastic Collision Expression}$$

$$v_{1'} = \frac{(m_1 - m_2)v_1 + 2m_2v_2}{m_1 + m_2} \quad \text{1-d Elastic Collision Expression}$$

$$v_{2'} - v_{1'} = -(v_2 - v_1) \quad v_{\text{rel}'} = -v_{\text{rel}} \quad \text{1-d Elastic Collision Expressions}$$

17 Rotational Kinematics

$$2\pi = 6.2831853\dots \quad \frac{1}{2\pi} = 0.15915494\dots$$

$$\frac{180^\circ}{\pi} = 57.295779\dots \approx 60^\circ \quad \frac{\pi}{180^\circ} = 0.017453292\dots \approx \frac{1}{60^\circ}$$

$$\theta = \frac{s}{r} \quad \omega = \frac{d\theta}{dt} = \frac{v}{r} \quad \alpha = \frac{d^2\theta}{dt^2} = \frac{d\omega}{dt} = \frac{a}{r} \quad f = \frac{\omega}{2\pi} \quad P = \frac{1}{f} = \frac{2\pi}{\omega}$$

$$\omega = \alpha t + \omega_0 \quad \Delta\theta = \frac{1}{2}\alpha t^2 + \omega_0 t \quad \omega^2 = \omega_0^2 + 2\alpha\Delta\theta$$

$$\Delta\theta = \frac{1}{2}(\omega_0 + \omega)t \quad \Delta\theta = -\frac{1}{2}\alpha t^2 + \omega t$$

18 Rotational Dynamics

$$\vec{L} = \vec{r} \times \vec{p} \quad \vec{\tau} = \vec{r} \times \vec{F} \quad \vec{\tau}_{\text{net}} = \frac{d\vec{L}}{dt}$$

$$L_z = RP_{xy} \sin \gamma_L \quad \tau_z = RF_{xy} \sin \gamma_\tau \quad L_z = I\omega \quad \tau_{z,\text{net}} = I\alpha$$

$$I = \sum_i m_i R_i^2 \quad I = \int R^2 \rho dV \quad I_{\text{parallel axis}} = I_{\text{cm}} + mR_{\text{cm}}^2 \quad I_z = I_x + I_y$$

$$I_{\text{cyl,shell,thin}} = MR^2 \quad I_{\text{cyl}} = \frac{1}{2}MR^2 \quad I_{\text{cyl,shell,thick}} = \frac{1}{2}M(R_1^2 + R_2^2)$$

$$I_{\text{rod,thin,cm}} = \frac{1}{12}ML^2 \quad I_{\text{sph,solid}} = \frac{2}{5}MR^2 \quad I_{\text{sph,shell,thin}} = \frac{2}{3}MR^2$$

$$a = \frac{g \sin \theta}{1 + I/(mr^2)}$$

$$KE_{\text{rot}} = \frac{1}{2}I\omega^2 \quad dW = \tau_z d\theta \quad P = \frac{dW}{dt} = \tau_z \omega$$

$$\Delta KE_{\text{rot}} = W_{\text{net}} = \int \tau_{z,\text{net}} d\theta \quad \Delta PE_{\text{rot}} = -W = -\int \tau_{z,\text{con}} d\theta$$

$$\Delta E_{\text{rot}} = KE_{\text{rot}} + \Delta PE_{\text{rot}} = W_{\text{non,rot}} \quad \Delta E = \Delta KE + KE_{\text{rot}} + \Delta PE = W_{\text{non}} + W_{\text{rot}}$$

19 Static Equilibrium

$$\vec{F}_{\text{ext,net}} = 0 \quad \vec{\tau}_{\text{ext,net}} = 0 \quad \vec{\tau}_{\text{ext,net}} = \tau'_{\text{ext,net}} \quad \text{if } F_{\text{ext,net}} = 0$$

$$0 = F_{\text{net } x} = \sum F_x \quad 0 = F_{\text{net } y} = \sum F_y \quad 0 = \tau_{\text{net}} = \sum \tau$$

20 Gravity

$$\vec{F}_{1 \text{ on } 2} = -\frac{Gm_1 m_2}{r_{12}^2} \hat{r}_{12} \quad \vec{g} = -\frac{GM}{r^2} \hat{r} \quad \oint \vec{g} \cdot d\vec{A} = -4\pi GM$$

$$PE = -\frac{Gm_1 m_2}{r_{12}} \quad V = -\frac{GM}{r} \quad v_{\text{escape}} = \sqrt{\frac{2GM}{r}} \quad v_{\text{orbit}} = \sqrt{\frac{GM}{r}}$$

$$P^2 = \left(\frac{4\pi^2}{GM}\right) r^3 \quad P = \left(\frac{2\pi}{\sqrt{GM}}\right) r^{3/2} \quad \frac{dA}{dt} = \frac{1}{2} r^2 \omega = \frac{L}{2m} = \text{Constant}$$

$$R_{\text{Earth,mean}} = 6371.0 \text{ km} \quad R_{\text{Earth,equatorial}} = 6378.1 \text{ km} \quad M_{\text{Earth}} = 5.9736 \times 10^{24} \text{ kg}$$

$$R_{\text{Earth mean orbital radius}} = 1.495978875 \times 10^{11} \text{ m} = 1.0000001124 \text{ AU} \approx 1.5 \times 10^{11} \text{ m} \approx 1 \text{ AU}$$

$$R_{\text{Sun,equatorial}} = 6.955 \times 10^8 \approx 109 \times R_{\text{Earth,equatorial}} \quad M_{\text{Sun}} = 1.9891 \times 10^{30} \text{ kg}$$

21 Fluids

$$\rho = \frac{\Delta m}{\Delta V} \quad p = \frac{F}{A} \quad p = p_0 + \rho g d_{\text{depth}}$$

$$\text{Pascal's principle} \quad p = p_{\text{ext}} - \rho g(y - y_{\text{ext}}) \quad \Delta p = \Delta p_{\text{ext}}$$

$$\text{Archimedes principle} \quad F_{\text{buoy}} = m_{\text{fluid dis}} g = V_{\text{fluid dis}} \rho_{\text{fluid}} g$$

$$\text{equation of continuity for ideal fluid} \quad R_V = Av = \text{Constant}$$

$$\text{Bernoulli's equation} \quad p + \frac{1}{2} \rho v^2 + \rho g y = \text{Constant}$$

22 Oscillation

$$P = f^{-1} \quad \omega = 2\pi f \quad F = -kx \quad U = \frac{1}{2} kx^2 \quad a(t) = -\frac{k}{m} x(t) = -\omega^2 x(t)$$

$$\omega = \sqrt{\frac{k}{m}} \quad P = 2\pi \sqrt{\frac{m}{k}} \quad x(t) = A \cos(\omega t) + B \sin(\omega t)$$

$$E_{\text{mec total}} = \frac{1}{2} m v_{\text{max}}^2 = \frac{1}{2} k x_{\text{max}}^2 = \frac{1}{2} m v^2 + \frac{1}{2} k x^2$$

$$P = 2\pi \sqrt{\frac{I}{mg\ell}} \quad P = 2\pi \sqrt{\frac{\ell}{g}}$$