

Intro Physics Semester I**Name:**

Homework 19: Thermodynamics II: Microscopic Treatment One or two full answer questions will be marked. There will also be a mark for completeness. Homeworks are due usually the day after the chapter they are for is finished. Solutions will be posted soon thereafter. The solutions are intended to be (but not necessarily are) super-perfect and often go beyond a fully correct answer.

Answer Table

	a	b	c	d	e		a	b	c	d	e
1.	O	O	O	O	O	31.	O	O	O	O	O
2.	O	O	O	O	O	32.	O	O	O	O	O
3.	O	O	O	O	O	33.	O	O	O	O	O
4.	O	O	O	O	O	34.	O	O	O	O	O
5.	O	O	O	O	O	35.	O	O	O	O	O
6.	O	O	O	O	O	36.	O	O	O	O	O
7.	O	O	O	O	O	37.	O	O	O	O	O
8.	O	O	O	O	O	38.	O	O	O	O	O
9.	O	O	O	O	O	39.	O	O	O	O	O
10.	O	O	O	O	O	40.	O	O	O	O	O
11.	O	O	O	O	O	41.	O	O	O	O	O
12.	O	O	O	O	O	42.	O	O	O	O	O
13.	O	O	O	O	O	43.	O	O	O	O	O
14.	O	O	O	O	O	44.	O	O	O	O	O
15.	O	O	O	O	O	45.	O	O	O	O	O
16.	O	O	O	O	O	46.	O	O	O	O	O
17.	O	O	O	O	O	47.	O	O	O	O	O
18.	O	O	O	O	O	48.	O	O	O	O	O
19.	O	O	O	O	O	49.	O	O	O	O	O
20.	O	O	O	O	O	50.	O	O	O	O	O
21.	O	O	O	O	O	51.	O	O	O	O	O
22.	O	O	O	O	O	52.	O	O	O	O	O
23.	O	O	O	O	O	53.	O	O	O	O	O
24.	O	O	O	O	O	54.	O	O	O	O	O
25.	O	O	O	O	O	55.	O	O	O	O	O
26.	O	O	O	O	O	56.	O	O	O	O	O
27.	O	O	O	O	O	57.	O	O	O	O	O
28.	O	O	O	O	O	58.	O	O	O	O	O
29.	O	O	O	O	O	59.	O	O	O	O	O
30.	O	O	O	O	O	60.	O	O	O	O	O

1. Classical thermodynamics as it existed in the 19th century avoided reference to microscopic entities like atoms and _____ since their existence was then considered problematic. If you make theories based on wrong premises, your theories are likely to be wrong. However, by the end of the 19th century evidence for atoms and _____ as real things was increasing and increasingly classical thermodynamics could be explained and extended by using the statistical behavior of the atoms and _____. The study of this statistical behavior is now called statistical mechanics. Although there is still some value in considering thermodynamics from a purely macroscopic point of view, it quickly becomes tedious to avoid referencing the microscopic realm since we know it is there and provides the basis for macroscopic thermodynamics. So it is physically right and pedagogically sound to start referencing the microscopic realm right from the start in studying thermodynamics.

- a) atoms
- b) molecules
- c) monads
- d) imperceptables
- e) little itty bitty things

Equation Sheet for Introductory Physics Calculus-Based

This equation sheet is intended for students writing tests or reviewing material. Therefore it is neither intended to be complete nor completely explicit. There are fewer symbols than variables, and so some symbols must be used for different things: context must distinguish.

The equations are mnemonic. Students are expected to understand how to interpret and use them.

1 Constants

$$c = 2.99792458 \times 10^8 \text{ m/s} \approx 2.998 \times 10^8 \text{ m/s} \approx 3 \times 10^8 \text{ m/s} \approx 1 \text{ lyr/yr} \approx 1 \text{ ft/ns} \quad \text{exact by definition}$$

$$e = 1.602176487(40) \times 10^{-19} \text{ C}$$

$$G = 6.67384(80) \times 10^{-11} \text{ N m}^2/\text{kg}^2 \quad (\text{2012, CODATA})$$

$$g = 9.8 \text{ m/s}^2 \quad \text{fiducial value}$$

$$k = \frac{1}{4\pi\varepsilon_0} = 8.987551787\dots \times 10^9 \approx 8.99 \times 10^9 \approx 10^{10} \text{ N m}^2/\text{C}^2 \text{ exact by definition}$$

$$k_{\text{Boltzmann}} = 1.3806504(24) \times 10^{-23} \text{ J/K} = 0.8617343(15) \times 10^{-4} \text{ eV/K} \approx 10^{-4} \text{ eV/K}$$

$$m_e = 9.10938215(45) \times 10^{-31} \text{ kg} = 0.510998910(13) \text{ MeV}$$

$$m_p = 1.672621637(83) \times 10^{-27} \text{ kg} = 938.272013(23) \text{ MeV}$$

$$\varepsilon_0 = \frac{1}{\mu_0 c^2} = 8.8541878176\dots \times 10^{-12} \text{ C}^2/(\text{N m}^2) \approx 10^{-11} \quad \text{vacuum permittivity (exact by definition)}$$

$$\mu_0 = 4\pi \times 10^{-7} \text{ N/A}^2 \quad \text{exact by definition}$$

2 Geometrical Formulae

$$C_{\text{cir}} = 2\pi r \quad A_{\text{cir}} = \pi r^2 \quad A_{\text{sph}} = 4\pi r^2 \quad V_{\text{sph}} = \frac{4}{3}\pi r^3$$

$$\Omega_{\text{sphere}} = 4\pi \quad d\Omega = \sin \theta \, d\theta \, d\phi$$

3 Trigonometry Formulae

$$\frac{x}{r} = \cos \theta \quad \frac{y}{r} = \sin \theta \quad \frac{y}{x} = \tan \theta = \frac{\sin \theta}{\cos \theta} \quad \cos^2 \theta + \sin^2 \theta = 1$$

$$\csc \theta = \frac{1}{\sin \theta} \quad \sec \theta = \frac{1}{\cos \theta} \quad \cot \theta = \frac{1}{\tan \theta}$$

$$c^2 = a^2 + b^2 \quad c = \sqrt{a^2 + b^2 - 2ab \cos \theta_c} \quad \frac{\sin \theta_a}{a} = \frac{\sin \theta_b}{b} = \frac{\sin \theta_c}{c}$$

$$f(\theta) = f(\theta + 360^\circ)$$

$$\sin(\theta + 180^\circ) = -\sin(\theta) \quad \cos(\theta + 180^\circ) = -\cos(\theta) \quad \tan(\theta + 180^\circ) = \tan(\theta)$$

$$\sin(-\theta) = -\sin(\theta) \quad \cos(-\theta) = \cos(\theta) \quad \tan(-\theta) = -\tan(\theta)$$

$$\sin(\theta + 90^\circ) = \cos(\theta) \quad \cos(\theta + 90^\circ) = -\sin(\theta) \quad \tan(\theta + 90^\circ) = -\tan(\theta)$$

$$\sin(180^\circ - \theta) = \sin(\theta) \quad \cos(180^\circ - \theta) = -\cos(\theta) \quad \tan(180^\circ - \theta) = -\tan(\theta)$$

$$\sin(90^\circ - \theta) = \cos(\theta) \quad \cos(90^\circ - \theta) = \sin(\theta) \quad \tan(90^\circ - \theta) = \frac{1}{\tan(\theta)} = \cot(\theta)$$

$$\sin(a + b) = \sin(a)\cos(b) + \cos(a)\sin(b) \quad \cos(a + b) = \cos(a)\cos(b) - \sin(a)\sin(b)$$

$$\sin(2a) = 2\sin(a)\cos(a) \quad \cos(2a) = \cos^2(a) - \sin^2(a)$$

$$\sin(a)\sin(b) = \frac{1}{2} [\cos(a - b) - \cos(a + b)] \quad \cos(a)\cos(b) = \frac{1}{2} [\cos(a - b) + \cos(a + b)]$$

$$\sin(a)\cos(b) = \frac{1}{2} [\sin(a - b) + \sin(a + b)]$$

$$\sin^2 \theta = \frac{1}{2}[1 - \cos(2\theta)] \quad \cos^2 \theta = \frac{1}{2}[1 + \cos(2\theta)] \quad \sin(a)\cos(a) = \frac{1}{2}\sin(2a)$$

$$\cos(x) - \cos(y) = -2 \sin\left(\frac{x+y}{2}\right) \sin\left(\frac{x-y}{2}\right)$$

$$\cos(x) + \cos(y) = 2 \cos\left(\frac{x+y}{2}\right) \cos\left(\frac{x-y}{2}\right)$$

$$\sin(x) + \sin(y) = 2 \sin\left(\frac{x+y}{2}\right) \cos\left(\frac{x-y}{2}\right)$$

4 Approximation Formulae

$$\frac{\Delta f}{\Delta x} \approx \frac{df}{dx} \quad \frac{1}{1-x} \approx 1+x : (x \ll 1)$$

$$\sin \theta \approx \theta \quad \tan \theta \approx \theta \quad \cos \theta \approx 1 - \frac{1}{2}\theta^2 \quad \text{all for } \theta \ll 1$$

5 Quadratic Formula

$$\text{If } 0 = ax^2 + bx + c, \quad \text{then } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = -\frac{b}{2a} \pm \sqrt{\left(\frac{b}{2a}\right)^2 - \frac{c}{a}}$$

6 Vector Formulae

$$a = |\vec{a}| = \sqrt{a_x^2 + a_y^2} \quad \theta = \tan^{-1} \left(\frac{a_y}{a_x} \right) + \pi? \quad \vec{a} + \vec{b} = (a_x + b_x, a_y + b_y)$$

$$a = |\vec{a}| = \sqrt{a_x^2 + a_y^2 + a_z^2} \quad \phi = \tan^{-1} \left(\frac{a_y}{a_x} \right) + \pi? \quad \theta = \cos^{-1} \left(\frac{a_z}{a} \right)$$

$$\vec{a} + \vec{b} = (a_x + b_x, a_y + b_y, a_z + b_z)$$

$$\vec{a} \cdot \vec{b} = ab \cos \theta = a_x b_x + a_y b_y + a_z b_z$$

$$\vec{c} = \vec{a} \times \vec{b} = ab \sin(\theta) \hat{c} = (a_y b_z - b_y a_z, a_z b_x - b_z a_x, a_x b_y - b_x a_y)$$

7 Differentiation and Integration Formulae

$$\frac{d(x^p)}{dx} = px^{p-1} \quad \text{except for } p = 0; \quad \frac{d(x^0)}{dx} = 0 \quad \frac{d(\ln|x|)}{dx} = \frac{1}{x}$$

Taylor's series

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} \frac{(x-x_0)^n}{n!} f^{(n)}(x_0) \\ &= f(x_0) + (x-x_0)f^{(1)}(x_0) + \frac{(x-x_0)^2}{2!}f^{(2)}(x_0) + \frac{(x-x_0)^3}{3!}f^{(3)}(x_0) + \dots \end{aligned}$$

$$\int_a^b f(x) dx = F(x)|_a^b = F(b) - F(a) \quad \text{where} \quad \frac{dF(x)}{dx} = f(x)$$

$$\int x^n dx = \frac{x^{n+1}}{n+1} \quad \text{except for } n = -1; \quad \int \frac{1}{x} dx = \ln|x|$$

8 One-Dimensional Kinematics

$$v_{\text{avg}} = \frac{\Delta x}{\Delta t} \quad v = \frac{dx}{dt} \quad a_{\text{avg}} = \frac{\Delta v}{\Delta t} \quad a = \frac{dv}{dt} = \frac{d^2x}{dt^2}$$

$$v = at + v_0 \quad x = \frac{1}{2}at^2 + v_0t + x_0 \quad v^2 = v_0^2 + 2a(x - x_0)$$

$$x = \frac{1}{2}(v_0 + v)t + x_0 \quad x = -\frac{1}{2}at^2 + vt + x_0 \quad g = 9.8 \text{ m/s}^2$$

$$x_{\text{rel}} = x_2 - x_1 \quad v_{\text{rel}} = v_2 - v_1 \quad a_{\text{rel}} = a_2 - a_1$$

$$x' = x - v_{\text{frame}}t \quad v' = v - v_{\text{frame}} \quad a' = a$$

9 Two- and Three-Dimensional Kinematics: General

$$\vec{v}_{\text{avg}} = \frac{\Delta \vec{r}}{\Delta t} \quad \vec{v} = \frac{d\vec{r}}{dt} \quad \vec{a}_{\text{avg}} = \frac{\Delta \vec{v}}{\Delta t} \quad \vec{a} = \frac{d\vec{v}}{dt} = \frac{d^2 \vec{r}}{dt^2}$$

10 Projectile Motion

$$x = v_{x,0}t \quad y = -\frac{1}{2}gt^2 + v_{y,0}t + y_0 \quad v_{x,0} = v_0 \cos \theta \quad v_{y,0} = v_0 \sin \theta$$

$$t = \frac{x}{v_{x,0}} = \frac{x}{v_0 \cos \theta} \quad y = y_0 + x \tan \theta - \frac{x^2 g}{2v_0^2 \cos^2 \theta}$$

$$x_{\text{for } y \text{ max}} = \frac{v_0^2 \sin \theta \cos \theta}{g} \quad y_{\text{max}} = y_0 + \frac{v_0^2 \sin^2 \theta}{2g}$$

$$x(y = y_0) = \frac{2v_0^2 \sin \theta \cos \theta}{g} = \frac{v_0^2 \sin(2\theta)}{g} \quad \theta_{\text{for max}} = \frac{\pi}{4} \quad x_{\text{max}}(y = y_0) = \frac{v_0^2}{g}$$

$$x(\theta = 0) = \pm v_0 \sqrt{\frac{2(y_0 - y)}{g}} \quad t(\theta = 0) = \sqrt{\frac{2(y_0 - y)}{g}}$$

11 Relative Motion

$$\vec{r} = \vec{r}_2 - \vec{r}_1 \quad \vec{v} = \vec{v}_2 - \vec{v}_1 \quad \vec{a} = \vec{a}_2 - \vec{a}_1$$

12 Polar Coordinate Motion and Uniform Circular Motion

$$\omega = \frac{d\theta}{dt} \quad \alpha = \frac{d\omega}{dt} = \frac{d^2\theta}{dt^2}$$

$$\vec{r} = r\hat{r} \quad \vec{v} = \frac{d\vec{r}}{dt} = \frac{dr}{dt}\hat{r} + r\omega\hat{\theta} \quad \vec{a} = \frac{d^2\vec{r}}{dt^2} = \left(\frac{d^2r}{dt^2} - r\omega^2 \right)\hat{r} + \left(r\alpha + 2\frac{dr}{dt}\omega \right)\hat{\theta}$$

$$\vec{v} = r\omega\hat{\theta} \quad v = r\omega \quad a_{\text{tan}} = r\alpha$$

$$\vec{a}_{\text{centripetal}} = -\frac{v^2}{r}\hat{r} = -r\omega^2\hat{r} \quad a_{\text{centripetal}} = \frac{v^2}{r} = r\omega^2 = v\omega$$

13 Very Basic Newtonian Physics

$$\vec{r}_{\text{cm}} = \frac{\sum_i m_i \vec{r}_i}{m_{\text{total}}} = \frac{\sum_{\text{sub}} m_{\text{sub}} \vec{r}_{\text{cm sub}}}{m_{\text{total}}} \quad \vec{v}_{\text{cm}} = \frac{\sum_i m_i \vec{v}_i}{m_{\text{total}}} \quad \vec{a}_{\text{cm}} = \frac{\sum_i m_i \vec{a}_i}{m_{\text{total}}}$$

$$\vec{r}_{\text{cm}} = \frac{\int_V \rho(\vec{r}) \vec{r} dV}{m_{\text{total}}}$$

$$\vec{F}_{\text{net}} = m\vec{a} \quad \vec{F}_{21} = -\vec{F}_{12} \quad F_g = mg \quad g = 9.8 \text{ m/s}^2$$

$$\vec{F}_{\text{normal}} = -\vec{F}_{\text{applied}} \quad F_{\text{linear}} = -kx$$

$$f_{\text{normal}} = \frac{T}{r} \quad T = T_0 - F_{\text{parallel}}(s) \quad T = T_0$$

$$F_{\text{f static}} = \min(F_{\text{applied}}, F_{\text{f static max}}) \quad F_{\text{f static max}} = \mu_{\text{static}} F_N \quad F_{\text{f kinetic}} = \mu_{\text{kinetic}} F_N$$

$$v_{\text{tangential}} = r\omega = r \frac{d\theta}{dt} \quad a_{\text{tangential}} = r\alpha = r \frac{d\omega}{dt} = r \frac{d^2\theta}{dt^2}$$

$$\vec{a}_{\text{centripetal}} = -\frac{v^2}{r} \hat{r} \quad \vec{F}_{\text{centripetal}} = -m \frac{v^2}{r} \hat{r}$$

$$F_{\text{drag,lin}} = bv \quad v_T = \frac{mg}{b} \quad \tau = \frac{v_T}{g} = \frac{m}{b} \quad v = v_T(1 - e^{-t/\tau})$$

$$F_{\text{drag,quad}} = bv^2 = \frac{1}{2} C \rho A v^2 \quad v_T = \sqrt{\frac{mg}{b}}$$

14 Energy and Work

$$dW = \vec{F} \cdot d\vec{s} \quad W = \int \vec{F} \cdot d\vec{s} \quad KE = \frac{1}{2}mv^2 \quad E_{\text{mechanical}} = KE + PE$$

$$P_{\text{avg}} = \frac{\Delta W}{\Delta t} \quad P = \frac{dW}{dt} \quad P = \vec{F} \cdot \vec{v}$$

$$\Delta KE = W_{\text{net}} \quad \Delta PE_{\text{of a conservative force}} = -W_{\text{by a conservative force}} \quad \Delta E = W_{\text{nonconservative}}$$

$$F = -\frac{dPE}{dx} \quad \vec{F} = -\nabla PE \quad PE = \frac{1}{2}kx^2 \quad PE = mgy$$

15 Momentum

$$\vec{F}_{\text{net}} = m\vec{a}_{\text{cm}} \quad \Delta KE_{\text{cm}} = W_{\text{net, external}} \quad \Delta E_{\text{cm}} = W_{\text{not}}$$

$$\vec{p} = m\vec{v} \quad \vec{F}_{\text{net}} = \frac{d\vec{p}}{dt} \quad \vec{F}_{\text{net}} = \frac{d\vec{p}_{\text{total}}}{dt}$$

$$m\vec{a}_{\text{cm}} = \vec{F}_{\text{net non-flux}} + (\vec{v}_{\text{flux}} - \vec{v}_{\text{cm}}) \frac{dm}{dt} = \vec{F}_{\text{net non-flux}} + \vec{v}_{\text{rel}} \frac{dm}{dt}$$

$$v = v_0 + v_{\text{ex}} \ln \left(\frac{m_0}{m} \right) \quad \text{rocket in free space}$$

16 Collisions

$$\vec{I} = \int_{\Delta t} \vec{F}(t) dt \quad \vec{F}_{\text{avg}} = \frac{\vec{I}}{\Delta t} \quad \Delta p = \vec{I}_{\text{net}}$$

$$\vec{p}_{1i} + \vec{p}_{2i} = \vec{p}_{1f} + \vec{p}_{2f} \quad \vec{v}_{\text{cm}} = \frac{\vec{p}_1 + \vec{p}_2}{m_{\text{total}}}$$

$$KE_{\text{total } f} = KE_{\text{total } i} \quad \text{1-d Elastic Collision Expression}$$

$$v_{1'} = \frac{(m_1 - m_2)v_1 + 2m_2v_2}{m_1 + m_2} \quad \text{1-d Elastic Collision Expression}$$

$$v_{2'} - v_{1'} = -(v_2 - v_1) \quad v_{\text{rel}'} = -v_{\text{rel}} \quad \text{1-d Elastic Collision Expressions}$$

17 Rotational Kinematics

$$2\pi = 6.2831853\dots \quad \frac{1}{2\pi} = 0.15915494\dots$$

$$\frac{180^\circ}{\pi} = 57.295779\dots \approx 60^\circ \quad \frac{\pi}{180^\circ} = 0.017453292\dots \approx \frac{1}{60^\circ}$$

$$\theta = \frac{s}{r} \quad \omega = \frac{d\theta}{dt} = \frac{v}{r} \quad \alpha = \frac{d^2\theta}{dt^2} = \frac{d\omega}{dt} = \frac{a}{r} \quad f = \frac{\omega}{2\pi} \quad P = \frac{1}{f} = \frac{2\pi}{\omega}$$

$$\omega = at + \omega_0 \quad \Delta\theta = \frac{1}{2}\alpha t^2 + \omega_0 t \quad \omega^2 = \omega_0^2 + 2\alpha\Delta\theta$$

$$\Delta\theta = \frac{1}{2}(\omega_0 + \omega)t \quad \Delta\theta = -\frac{1}{2}\alpha t^2 + \omega t$$

18 Rotational Dynamics

$$\vec{L} = \vec{r} \times \vec{p} \quad \vec{\tau} = \vec{r} \times \vec{F} \quad \vec{\tau}_{\text{net}} = \frac{d\vec{L}}{dt}$$

$$L_z = RP_{xy} \sin \gamma_L \quad \tau_z = RF_{xy} \sin \gamma_\tau \quad L_z = I\omega \quad \tau_{z,\text{net}} = I\alpha$$

$$I = \sum_i m_i R_i^2 \quad I = \int R^2 \rho \, dV \quad I_{\text{parallel axis}} = I_{\text{cm}} + mR_{\text{cm}}^2 \quad I_z = I_x + I_y$$

$$I_{\text{cyl,shell,thin}} = MR^2 \quad I_{\text{cyl}} = \frac{1}{2}MR^2 \quad I_{\text{cyl,shell,thick}} = \frac{1}{2}M(R_1^2 + R_2^2)$$

$$I_{\text{rod,thin,cm}} = \frac{1}{12}ML^2 \quad I_{\text{sph,solid}} = \frac{2}{5}MR^2 \quad I_{\text{sph,shell,thin}} = \frac{2}{3}MR^2$$

$$a = \frac{g \sin \theta}{1 + I/(mr^2)}$$

$$KE_{\text{rot}} = \frac{1}{2}I\omega^2 \quad dW = \tau_z \, d\theta \quad P = \frac{dW}{dt} = \tau_z \omega$$

$$\Delta KE_{\text{rot}} = W_{\text{net}} = \int \tau_{z,\text{net}} \, d\theta \quad \Delta PE_{\text{rot}} = -W = - \int \tau_{z,\text{con}} \, d\theta$$

$$\Delta E_{\text{rot}} = KE_{\text{rot}} + \Delta PE_{\text{rot}} = W_{\text{non,rot}} \quad \Delta E = \Delta KE + KE_{\text{rot}} + \Delta PE = W_{\text{non}} + W_{\text{rot}}$$

19 Static Equilibrium

$$\vec{F}_{\text{ext,net}} = 0 \quad \vec{\tau}_{\text{ext,net}} = 0 \quad \vec{\tau}_{\text{ext,net}} = \tau'_{\text{ext,net}} \quad \text{if } F_{\text{ext,net}} = 0$$

$$0 = F_{\text{net } x} = \sum F_x \quad 0 = F_{\text{net } y} = \sum F_y \quad 0 = \tau_{\text{net}} = \sum \tau$$

20 Gravity

$$\vec{F}_{1 \text{ on } 2} = -\frac{Gm_1m_2}{r_{12}^2} \hat{r}_{12} \quad \vec{g} = -\frac{GM}{r^2} \hat{r} \quad \oint \vec{g} \cdot d\vec{A} = -4\pi GM$$

$$PE = -\frac{Gm_1m_2}{r_{12}} \quad V = -\frac{GM}{r} \quad v_{\text{escape}} = \sqrt{\frac{2GM}{r}} \quad v_{\text{orbit}} = \sqrt{\frac{GM}{r}}$$

$$P^2 = \left(\frac{4\pi^2}{GM}\right) r^3 \quad P = \left(\frac{2\pi}{\sqrt{GM}}\right) r^{3/2} \quad \frac{dA}{dt} = \frac{1}{2}r^2\omega = \frac{L}{2m} = \text{Constant}$$

$$R_{\text{Earth,mean}} = 6371.0 \text{ km} \quad R_{\text{Earth,equatorial}} = 6378.1 \text{ km} \quad M_{\text{Earth}} = 5.9736 \times 10^{24} \text{ kg}$$

$$R_{\text{Earth mean orbital radius}} = 1.495978875 \times 10^{11} \text{ m} = 1.0000001124 \text{ AU} \approx 1.5 \times 10^{11} \text{ m} \approx 1 \text{ AU}$$

$$R_{\text{Sun,equatorial}} = 6.955 \times 10^8 \approx 109 \times R_{\text{Earth,equatorial}} \quad M_{\text{Sun}} = 1.9891 \times 10^{30} \text{ kg}$$

21 Fluids

$$\rho = \frac{\Delta m}{\Delta V} \quad p = \frac{F}{A} \quad p = p_0 + \rho g d_{\text{depth}}$$

$$\text{Pascal's principle} \quad p = p_{\text{ext}} - \rho g(y - y_{\text{ext}}) \quad \Delta p = \Delta p_{\text{ext}}$$

$$\text{Archimedes principle} \quad F_{\text{buoy}} = m_{\text{fluid dis}} g = V_{\text{fluid dis}} \rho_{\text{fluid}} g$$

$$\text{equation of continuity for ideal fluid} \quad R_V = Av = \text{Constant}$$

$$\text{Bernoulli's equation} \quad p + \frac{1}{2} \rho v^2 + \rho g y = \text{Constant}$$

22 Oscillation

$$P = f^{-1} \quad \omega = 2\pi f \quad F = -kx \quad PE = \frac{1}{2} kx^2 \quad a(t) = -\frac{k}{m}x(t) = -\omega^2 x(t)$$

$$\omega = \sqrt{\frac{k}{m}} \quad P = 2\pi \sqrt{\frac{m}{k}} \quad x(t) = A \cos(\omega t) + B \sin(\omega t)$$

$$E_{\text{mec total}} = \frac{1}{2}mv_{\text{max}}^2 = \frac{1}{2}kx_{\text{max}}^2 = \frac{1}{2}mv^2 + \frac{1}{2}kx^2$$

$$P = 2\pi \sqrt{\frac{I}{mgr}} \quad P = 2\pi \sqrt{\frac{r}{g}}$$

23 Waves

$$\frac{d^2y}{dx^2} = \frac{1}{v^2} \frac{d^2y}{dt^2} \quad v = \sqrt{\frac{F_{\text{T}}}{\mu}} \quad y = f(x \mp vt)$$

$$y = y_{\text{max}} \sin[k(x \mp vt)] = y_{\text{max}} \sin(kx \mp \omega t)$$

$$\text{Period} = \frac{1}{f} \quad k = \frac{2\pi}{\lambda} \quad v = f\lambda = \frac{\omega}{k} \quad P \propto y_{\text{max}}^2$$

$$y = 2y_{\max} \sin(kx) \cos(\omega t) \quad n = \frac{L}{\lambda/2} \quad L = n \frac{\lambda}{2} \quad \lambda = \frac{2L}{n} \quad f = n \frac{v}{2L}$$

$$v = \sqrt{\left(\frac{\partial P}{\partial \rho}\right)_S} \quad n\lambda = d \sin(\theta) \quad \left(n + \frac{1}{2}\right)\lambda = d \sin(\theta)$$

$$I = \frac{P}{4\pi r^2} \quad \beta = (10 \text{ dB}) \times \log \left(\frac{I}{I_0} \right)$$

$$f = n \frac{v}{4L} : n = 1, 3, 5, \dots \quad f_{\text{medium}} = \frac{f_0}{1 - v_0/v_{\text{medium}}}$$

$$f' = f \left(1 - \frac{v'}{v} \right) \quad f = \frac{f'}{1 - v'/v}$$

24 Thermodynamics

$$dE = dQ - dW = T dS - p dV$$

$$T_{\text{K}} = T_{\text{C}} + 273.15 \text{ K} \quad T_{\text{F}} = 1.8 \times T_{\text{C}} + 32^{\circ}\text{F}$$

$$Q = mC\Delta T \quad Q = mL$$

$$PV = NkT \quad P = \frac{2}{3} \frac{N}{V} KE_{\text{avg}} = \frac{2}{3} \frac{N}{V} \left(\frac{1}{2} mv_{\text{RMS}}^2 \right)$$

$$v_{\text{RMS}} = \sqrt{\frac{3kT}{m}} = 2735.51 \dots \times \sqrt{\frac{T/300}{A}}$$

$$PV^\gamma = \text{constant} \quad 1 < \gamma \leq \frac{5}{3} \quad v_{\text{sound}} = \sqrt{\frac{B}{\rho}} = \sqrt{\frac{-V(\partial P/\partial V)_S}{m(N/V)}} = \sqrt{\frac{\gamma kT}{m}}$$

$$\varepsilon = \frac{W}{Q_{\text{H}}} = \frac{Q_{\text{H}} - Q_{\text{C}}}{W} = 1 - \frac{Q_{\text{C}}}{Q_{\text{H}}}$$

$$\eta_{\text{heating}} = \frac{Q_{\text{H}}}{W} = \frac{Q_{\text{H}}}{Q_{\text{H}} - Q_{\text{C}}} = \frac{1}{1 - Q_{\text{C}}/Q_{\text{H}}} = \frac{1}{\varepsilon} \quad \eta_{\text{cooling}} = \frac{Q_{\text{C}}}{W} = \frac{Q_{\text{H}} - W}{W} = \frac{1}{\varepsilon} - 1 = \eta_{\text{heating}} - 1$$

$$\varepsilon_{\text{Carnot}} = 1 - \frac{T_{\text{C}}}{T_{\text{H}}} \quad \eta_{\text{heating,Carnot}} = \frac{1}{1 - T_{\text{C}}/T_{\text{H}}} \quad \eta_{\text{cooling,Carnot}} = \frac{T_{\text{C}}/T_{\text{H}}}{1 - T_{\text{C}}/T_{\text{H}}}$$